

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/341272117>

Dietary pattern recognition on Twitter: a case example of before, during, and after four natural disasters

Article in *Natural Hazards* · May 2020

DOI: 10.1007/s11069-020-04024-6

CITATIONS

0

READS

11

4 authors, including:



Amir Karami

University of South Carolina

40 PUBLICATIONS 482 CITATIONS

[SEE PROFILE](#)



Courtney Monroe

University of South Carolina

18 PUBLICATIONS 124 CITATIONS

[SEE PROFILE](#)



Heather M. Brandt

University of South Carolina

183 PUBLICATIONS 3,793 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Create new project "The ImPACT Feasibility Study" [View project](#)



SCCDCN-II [View project](#)



Dietary pattern recognition on Twitter: a case example of before, during, and after four natural disasters

Gabrielle Turner-McGrievy¹ · Amir Karami² · Courtney Monroe¹ · Heather M. Brandt¹

Received: 3 July 2019 / Accepted: 25 April 2020
© Springer Nature B.V. 2020

Abstract

Little is known about what foods/beverages (F&B) are common during natural disasters. The goal of this study was to track high-frequency F&B mentions during four hurricanes affecting the coast of South Carolina for quantifying dietary patterns in Twitter. A listing of common F&B ($n=173$) was created from the top food sources of energy, fat, protein, and carbohydrate in the USA. A sampling of > 500,000 tweets containing hashtag names (e.g., #HurricaneFlorence) or actual names (e.g., “Hurricane Florence”) of the four hurricanes was collected using Crimson Hexagon. ANOVA was used to examine differences in number of mentions in each food group pre- (6 days before), during (48 h of the hurricane), and post-hurricane (6 days after). Descriptive statistics were used to examine the most frequently mentioned F&B (threshold defined as ≥ 4 mentions/day for each F&B item or 10% of the foods mentioned) and whether F&B were top sources of energy/macronutrients. More than 5000 mentions of F&B were collected in our sample. Grains were the most frequently mentioned food group pre-hurricane, and dairy was most frequently mentioned during the hurricanes. The top five most commonly mentioned F&B overall were milk ($n=517$), pizza ($n=511$), turkey ($n=425$), oranges ($n=384$), and waffles ($n=346$). Foods mentioned were commonly energy and protein dense. Five foods (pizza, waffles, milk, rolls, and bread) were categorized as a top contributor across energy and all three macronutrients. Social media may be a unique way to detect dietary patterns and help inform public health social media campaigns to advise people about stocking up on healthy, non-perishable foods ahead of natural disasters.

✉ Gabrielle Turner-McGrievy
brie@sc.edu

Amir Karami
karami@sc.edu

Courtney Monroe
cmonroe@mailbox.sc.edu

Heather M. Brandt
hbrandt@sc.edu

¹ Arnold School of Public Health, University of South Carolina, 915 Greene Street, Columbia, SC 29208, USA

² School of Library and Information Science, University of South Carolina, 1501 Greene Street, Columbia, SC 29208, USA

Keywords Diet · Diet patterns · Twitter · Natural disaster · Social media

1 Introduction

Every day, millions of users post messages on social media platforms, such as Twitter, Facebook, or Instagram (Smith 2019). While Facebook remains the most popular social media platform in the USA, all social media platforms have seen increased usage over the past decade, including Twitter (Smith and Anderson 2018). Nearly a quarter of all internet users are also Twitter users (Smith and Anderson 2018). Twitter is an online social networking platform that lets users submit posts that are 280 characters or less. Twitter, which has been in existence since 2006, has emerged as a powerful research tool for both behavioral interventions and for observational data collection of human behaviors (Sinnenberg et al. 2017). This research has included using Twitter as a source of data and using Twitter to recruit and/or intervene on participants (Sinnenberg et al. 2017).

The purpose of this study was to utilize Twitter as a way to assess dietary patterns before, during, and after four natural disasters, including Hurricane Matthew 2016, Hurricane Irma 2017, Hurricane Florence 2018, and Hurricane Michael 2018 that affected South Carolina and surrounding areas. The search was limited to these four major storms affecting a similar geographical area in order to control for potential regional food preferences. These storms caused significant financial and structural damage. For example, Hurricane Florence caused an estimated \$24 billion in damages to Virginia, North Carolina, and South Carolina when it made landfall on September 14, 2018 (Duncan 2019). Because of the widespread impact of these hurricanes and the need for preparation efforts ahead of landfall, hurricanes represent an opportune event to study the use of Twitter to detect dietary patterns and food mentions before, during, and after a natural disaster.

One area of Twitter-based research is characterized by health-related and behavioral topics, including nutrition and other food-related subjects (Sinnenberg et al. 2017). Twitter has been used to identify food-borne illness reports and then encourage individuals to report these illnesses (Harris et al. 2017). Twitter has also been used to examine characteristics of the neighborhood food environment (Nguyen et al. 2017), the relationship between obesity rates and geo-coded Twitter posts related to food intake (Gore et al. 2015), the context in which food choices are made (Vidal et al. 2015), the topics posted related to diet, diabetes, exercise, and obesity (Karami et al. 2018), or how certain food topics are discussed (Kuttschreuter and Hilverda 2019). In addition, researchers have examined what factors make Twitter messages about healthy eating more likely to be re-tweeted and shared across social networks (Zhou et al. 2018). Twitter has also been used to help users track food intake and examine the relationship between dietary and behavioral factors (Hingle et al. 2013).

Another area of Twitter-based research focuses on the use of the platform during natural disasters (Karami et al. 2020; Guan and Chen 2014). Twitter was widely used as a way to disseminate crisis information during Hurricane Sandy (Wang and Zhuang 2017) and was one of the most discussed topics on Twitter, with 34% of all posts to Twitter during the 3 days around the storm being related to the hurricane (Heimlich 2012). In addition, Twitter was used to communicate time-sensitive and critical information during a historic flood that happened in the Midlands of South Carolina in 2015 (Brandt et al. in press; Karami et al. 2019). Communities affected by natural disasters have come to rely on social media as a way to exchange information about rescue and recovery efforts (Finch et al.

2016; Guskin and Hirtline 2012; Wang et al. 2016; Xiao et al. 2015). In addition, social media has been used to help victims, mobilize volunteers, and allow community members to communicate efficiently during both natural disasters and human-made or terrorist situations (Bennett 2014; Scott and Errett 2018; Sutton et al. 2015). It can also be used to spread misinformation during times of crisis (Wang and Zhuang 2018).

There are important food and beverage (F&B) guidelines that should be followed before, during, and after a natural disaster, such as a hurricane. The Centers for Disease Control and Prevention (CDC) recommends that people have five gallons of water per person on hand in preparation of a hurricane (CDC 2017). Specifically related to food, the CDC recommends having non-perishable food on hand, that has a long shelf life, requires little to no cooking or refrigeration and is low salt and not spicy (CDC 2019). Alcohol intake may increase during stressful events, like hurricanes, and because alcohol impairs judgement, the CDC recommends alcohol be avoided during natural disasters (CDC 2014). The previous research has shown that alcohol usage also tends to increase post-hurricane among those who have survived a hurricane (Flory et al. 2009). News articles in hurricane-prone areas have urged people not to overindulge in alcohol during storms, since hurricane parties (a practice of hosting parties, often involving alcohol consumption during a hurricane) are common (Dedaj 2017). In addition, popular press articles have urged people to stock up on healthy food to have on hand for a hurricane (Laseter 2018).

This study had two primary objectives. The first objective of this study was to develop a way to quantify food patterns in order to assess which foods were mentioned on Twitter and which are major contributors of energy and macronutrients. The second objective was to examine which foods and associated sources of energy and macronutrients were most frequently mentioned before, during, and after four natural disasters.

2 Materials and methods

Figure 1 details how F&B were collected, organized, and analyzed. The first step in this project was to develop a list of common food groups, foods that could be categorized into each food group, and then categorize those foods into the top F&B sources of energy, protein, fat, and carbohydrate in the USA. In order to develop this list, information detailing the top F&B sources of energy and macronutrients using National Health and Nutrition Examination Survey (NHANES) data for adults aged 19 years and older was used (O'Neil et al. 2012). In order to categorize mentions of F&B on Twitter into common food groups, the 10 food groups listed in the O'Neil et al. (2012) article were used. The 10 food groups




Fig. 1 How foods and beverages were collected, organized, and analyzed

were grain products; vegetables; fruit; dairy products; meat, poultry, and fish; eggs, legumes, nuts, and seeds; fats and oils; desserts and sweets; beverages; and other foods (e.g., condiments, broth, salt) (O’Neil et al. 2012). Next, the top 15 F&B sources of energy and macronutrients were used for analysis (with the exception of protein, which only had 12 top sources listed) (O’Neil et al. 2012). In order to ensure a variety of foods mentioned for a particular food group was captured, a list of other common names for the foods listed was developed. For example, a list of other common words was developed for broad food terms listed in the article like beef (e.g., created additional Twitter search terms such as hamburger, burger, and steak), poultry (e.g., chicken and turkey as additional terms), and pastry (e.g., donut, doughnut, and Danish as additional terms). In particular, three terms were very broad food categories (fruit, fish, and alcohol). In order to develop a comprehensive list of terms that could be used for those food categories, we conducted a Google web search for “top types of (fruit, fish, or alcohol) sold in the US.” Those web search results helped to complete a more comprehensive list of potential terms (e.g., instead of just searching for fruit, we also included bananas, apples, grapes, strawberries, oranges, etc.). This resulted in a list of common F&B items consumed in the USA that were top contributors to energy, protein, fat, and/or carbohydrate.

In order to examine food mentions during the hurricanes, a sampling of 506,620 tweets were collected during the 6 days before and after and 2 days during the hurricanes using a Twitter data provider, Crimson Hexagon (<https://www.crimsonhexagon.com/>) (see Table 1). For example, Hurricane Florence made landfall on September 14, 2018, so for that hurricane, pre-hurricane tweets were collected from September 8–13, during hurricane tweets were from September 14–15, and post-hurricane tweets were from September 16–21. Tweets needed to contain either the hashtag name for the hurricane (e.g., #HurricaneFlorence) or could contain the name of the hurricane (e.g., “Hurricane Florence”). A list of 173 F&B terms and their food group categories was created, along with possible plural versions of any food name that could commonly be plural (e.g., both pizza and pizzas were searched). Next, those tweets that contained a mention of one of the 173 F&B terms were collected, and frequency of tweets by date for each food term was categorized. This process was conducted using R Statistical Software (RCoreTeam 2014). Total posts during each time point (overall, pre-, during, and post-hurricane) were examined. In order to assess F&B and associated sources of energy and macronutrients with higher levels of mentions, a threshold was defined as ≥ 4 mentions/day for each F&B item or 10% of the

Table 1 Number of total tweets for each hurricane with either the hashtag name (e.g., #HurricaneMatthew) or regular name (e.g., Hurricane Matthew) the 6 days before, the 2 days during, and the 6 days after each hurricane and for overall

Hurricane	Number of Tweets			
	Before	During	After	Total
Matthew, 2016	59,442 (October 2–7)	19,433 (October 8–9)	49,529 (October 10–15)	128,404
Irma, 2017	52,244 (September 4–9)	17,281 (September 10–11)	54,180 (September 12–17)	123,705
Florence, 2018	56,071 (September 8–13)	17,419 (September 14–15)	54,599 (September 16–21)	128,089
Michael, 2018	49,699 (October 4–9)	20,000 (October 10–11)	56,723 (October 12–17)	126,422
Total	217,456	74,133	215,031	506,620

foods mentioned. This approach allowed for a focus on the most commonly mentioned foods and to attempt to limit counting mentions of foods that might have been infrequently tweeted, but were unrelated to food (e.g., I hope this weather “rolls” through quickly). Table 2 provides a list of example search terms for the ten food and beverage (F&B) categories along with example tweets.

2.1 Statistical methods

A one-way ANOVA with Tukey’s follow-up post hoc test was conducted in order to examine differences in mean mentions of foods and beverages in category (\pm SD) examining total posts and posts made pre-, during, and post-hurricane, regardless of threshold. A P value of <0.05 was used to indicate statistical significance. Descriptive statistics were used to examine the top most frequently mentioned foods and beverages (that met the ≥ 4 mentions/day threshold) and whether those foods and beverages were top sources of energy or macronutrients.

3 Results

In our sample, 5219 mentions of foods and beverages were detected. Table 3 shows the total mean (\pm SD) mentions of foods and beverages by food group category over the entire study period (14 days) as well as pre-hurricane (6 days before), during hurricane (48 h of the hurricane), and post-hurricane (6 days after the hurricane). While there were no significant differences among the F&B categories for total mentions ($P=0.12$) or mentions after the hurricanes ($P=0.50$), there were significant differences in food group categories tweeted pre- and during the hurricanes. Pre-hurricane, there were significantly more posts categorized as grains than those categorized as vegetables, meat/poultry/fish, eggs/legumes/nuts/seeds, or other (P ’s all <0.05). During the hurricanes, there were significantly more posts categorized as dairy than all other food groups ($P<0.05$). Figure 2 details the frequency of mentions within each food group category by day examined (each of the 6 days before and after the hurricanes and the 2 days during).

Considering the threshold of examining only those F&B that had ≥ 4 mentions per day, there were 13 F&B mentioned pre-hurricane, 18 during the hurricane, and 21 post-hurricane. Number of foods mentioned within each F&B group category was also examined. Three out of the 10 categories did not have F&B mentioned over the course of the entire examined period that met the ≥ 4 mentions per day threshold (vegetables, desserts and sweets, and items categorized as other). Another two food group categories had only one food or beverage item mentioned within that category. For dairy, only milk was mentioned ($n=517$) with no mentions of other dairy foods, such as cheese, and for fats and oils, only oil was mentioned ($n=135$). Of the remaining five food group categories, grains had five commonly tweeted foods (pizza, waffles, spaghetti, rolls, bread, $n=1735$), fruit had two commonly tweeted foods (apples and oranges, $n=505$), meat/poultry/fish had five commonly tweeted foods (catfish, chicken, fish, poultry, and turkey, $n=880$), eggs/legumes/nuts/seeds had two commonly tweeted foods (eggs and peanuts, $n=237$), and beverages had four commonly tweeted items (beer, coffee, alcohol, and wine, $n=515$). Figure 3 displays a word cloud (Heimerl et al. 2014) of the most commonly tweeted F&B overall. The top five most commonly mentioned F&B overall were milk ($n=517$), pizza ($n=511$), turkey ($n=425$), oranges ($n=384$), and waffles ($n=346$). Pre-hurricane, the top five foods

Table 2 Specific search terms and example tweets for the ten food and beverage (F&B) categories

	Example F&B search terms				Example tweets		
Grains	Flour	Rice	Pancakes	Pretzels	When a tree falls on your fence in the backyard because of #HurricaneMichael, you forgo cooking dinner and order pizza. Happy Thursday!		
	Bran	Pasta	Cracker	Chips			
	Bread	Biscuit	Crackers	Pizzas			
	Rolls	Biscuits	Popcorn	Pizzas			
	Cereal	Pancake	Pretzel	Cornbread			
Vegetables	Potato	Tomato	Lima bean	Pickles	Wax Beans	We cannot run out of potato salad again. #FEMA #HurricaneIrma #TheRingOfFire	
	Potatoes	Lettuce	Lima beans	Vegetable	Yellow bean		
	Broccoli	Green bean	Olive	Vegetables	Yellow beans		
	Spinach	Green beans	Olives	French fries	String bean		
	Squash	Corn	Pickle	Wax bean	String beans		
	Tomatoes	Peas					
	Juice	Apples	Watermelon	Peaches	Cherry		Thought I'd stop at the grocery store and pick up some apples on my way home... Nope. (#HurricaneMatthew)
	Fruit juice	Grapes	Watermelons	Cantaloupe	Cherries		
	Fruit	Grapes	Lemon	Cantaloupes	Pear		
	Smoothie	Strawberry	Lemons	Avocado	Pears		
Banana	Strawberries	Blueberry	Avocados	Lime			
Dairy products	Bananas	Orange	Blueberries	Pineapple	Limes	Hurricane shopping in SC. Where not only do people snatch up bread, milk, eggs, and water, but spaghetti and Vienna sausages. #HurricaneFlorence #southcarolina #hurricanesupplies #groceries #chs #Florence #hurricaneSeason2018 #HurricaneSeason #EastCoast #Atlantic	
	Apple	Oranges	Peach	Pineapples			
	Milk	Cheese	Chocolate milk	Cheez whiz	Velveeta		
	Yogurt	Cheeses					

Table 2 (continued)

Example F&B search terms				Example tweets	
Meat, Poultry, Fish	Beef	Frankfurters	Steak	Shrimp	Salami
	Lamb	Sausage	Steaks	Turkey	Pollock
	Veal	Luncheon Meat	Liver	Salmon	Crab
	Pork	Poultry	Hot dog	Lobster	Pepperoni
	Ham	Fish	Baloney	Pastrami	Catfish
Eggs, legumes, nuts, and seeds	Bacon	Shellfish	Bologna	Tilapia	Clams
	Organ meat	Burger	Chicken	Scallops	Cod
	Organ Meats	Hamburger	Tuna		
	Egg	Tofu	Nuts	Almonds	Seeds
	Eggs	Soy	Peanut	Almond butter	Sunflower Seed
Fats and oils	Legume	Tempeh	Peanuts	Cashew	Sunflower Seeds
	Legumes	Edamame	Peanut butter	Cashews	Flaxseed
	Bean	Veggie burger	Almond	Seed	Chia seed
	Beans	Nut			
	Margarine	Butter	Salad dressing	Mayonnaise	Oil
Desserts and sweets	Cake	Quick bread	Pies	Banana bread	Danishes
	Cakes	Pastry	Milk desserts	Donut	Ice cream
	Cookie	Pastries	Candy	Doughnut	Chocolate
	Cookies	Pie	Candies	Danish	Frozen yogurt
Beverages	Fruit drink	Kool-aid	Beer	Vodka	Whiskey
	Ades	Gatorade	Beers	Tequila	Cognac
	Coffee	Powerade	Wine	Rum	Bourbon
	Tea	Alcohol	Liquor	Gin	
Other foods	Meal-replacements	Condiments	Artificial Sweeteners	Marinara	Nutri-sweet
	Soup	Sauce	Salt	Splenda	Salsa
	Soups	Sauces	Slim-fast	Ketchup	Stevia
	Broth	Whey	Mustard	Barbecue	

Ugh, here we go again...people stupid insane raiding stores for milk, eggs, bread, water, and gasoline. Went to get hamburger buns for lunch and bacon and eggs for breakfast then refill lawnmower can—I found one gas station half sold out and WalMart insane. #HurricaneFlorence

Bought bread, tuna, and peanut butter. Better safe than sorry ____ #HurricaneFlorence #chswx

*Covers house in coconut oil #HurricaneMatthew

This just in the cookies i made are delicious #HurricaneMichael

Went to check on my local grocery stores in #ClaytonNC. Out of all water, sweet tea and most juice. #hurricaneflorence #ncwx

RT @jacquittagain Everyone: * grabs bread and milk for hurricane prep* Me: *grabs chips and salsa* #Hurricane-Florence

Table 3 Mean total mentions on Twitter before, during, and after Hurricanes Florence, Michael, Irma, and Matthew within ten food and beverage (F&B) categories

	Mean total mentions within each F&B category (\pm SD) examining the 6 days before the hurricanes	Mean total mentions within each food and beverage category (\pm SD) examining the 2 days of the hurricanes	Mean total mentions within each food and beverage category (\pm SD) examining the 6 days after the hurricanes	Mean total mentions within each food and beverage category (\pm SD) examining all posts pre, during, and post-hurricane
ANOVA model significance	$F = 2.79$; $P = 0.005$	$F = 3.11$; $P = 0.002$	$F = 0.93$; $P = 0.50$	$F = 1.61$; $P = 0.12$
Grains	27.4 \pm 46.5	7.8 \pm 14.9 ^b	41.1 \pm 105.2	76.3 \pm 138.5
Vegetables	1.1 \pm 2.3 ^a	0.4 \pm 0.7 ^b	3.7 \pm 7.6	5.1 \pm 8.7
Fruit	9.5 \pm 25.7	6.5 \pm 17.8 ^b	16.5 \pm 42.2	32.2 \pm 85.1
Dairy products	9.7 \pm 19.5	73.8 \pm 179.9	5.0 \pm 10.8	88.5 \pm 210.0
Meat, poultry, fish	3.7 \pm 6.5 ^a	3.3 \pm 7.0 ^b	24.2 \pm 65.7	31.2 \pm 76.2
Eggs, legumes, nuts, and seeds	3.7 \pm 6.5 ^a	0.7 \pm 1.2 ^b	11.5 \pm 35.5	15.8 \pm 41.1
Fats and oils	16.6 \pm 32.3	1.6 \pm 3.1 ^b	15.4 \pm 23.8	33.6 \pm 58.5
Desserts and sweets	4.3 \pm 6.4	0.7 \pm 1.6 ^b	6.6 \pm 8.4	11.6 \pm 13.5
Beverages	15.7 \pm 26.1 ^a	4.4 \pm 8.8 ^b	14.6 \pm 27.3	34.7 \pm 54.0
Other foods	2.1 \pm 5.0	0.7 \pm 2.3 ^b	3.3 \pm 6.7	6.1 \pm 12.5

^aSignificantly different from grains ($P < 0.05$)^bSignificantly different from dairy ($P < 0.05$)




Fig. 2 Frequency of mentions within each food group category by day examined

Fig. 3 Word cloud of the most commonly tweeted foods and beverages overall



mentioned were mostly grain-based foods: waffles ($n=142$), oranges ($n=117$), spaghetti ($n=117$), bread ($n=96$), and rolls ($n=79$). During the hurricanes, the top five foods mentioned were milk ($n=441$), oranges ($n=75$), pizza ($n=46$), waffles ($n=43$), and apples ($n=38$). Top five food mentions post-hurricane were pizza ($n=436$), turkey ($n=377$), oranges ($n=192$), waffles ($n=161$), and peanuts ($n=158$).

Lastly, we examined whether the F&B meeting the ≥ 4 mentions per day threshold were major contributors to energy, protein, fat, and/or carbohydrate intake in the USA. Of the 20 F&B mentioned, 60% were top sources of energy, 60% were top sources of protein, 45% were top sources of fat, and 40% were top sources of carbohydrate. Five foods mentioned (pizza, waffles, milk, rolls, and bread) were categorized as a top contributor across energy and all three macronutrient categories; three foods mentioned (turkey, chicken, and poultry) were common across energy and two macronutrient categories (fat and protein); one food mentioned (oil) was common across energy and one macronutrient category (fat); ten foods mentioned were a contributor to just one energy or macronutrient category (energy: alcohol, beer, wine; protein: peanuts, fish, catfish, eggs; carbohydrate: oranges, apples, spaghetti). One F&B (coffee) was not a major contributor to US intakes of energy or macronutrients.

4 Discussion

When a natural disaster, such as a hurricane, is predicted to occur, it is recommended to have a supply of non-perishable, healthy food on hand in case there is a loss of electricity or an inability to leave the house to shop for food (CDC 2017). Food is a common item people cite when preparing for a natural disaster, more so than ensuring adequate water, flashlights, medications, and cash (Wang 2017). Most households have at least a three-day supply of food on hand (Der-Martirosian et al. 2014). However, very little is known about what foods or beverages people tend to purchase or consume before, during, or after natural disasters. Twitter provides a unique opportunity to examine what F&B are commonly mentioned during natural disasters. Unique to this study, assessment of food groups and major contributors of energy and macronutrients was conducted in order to examine broader dietary patterns during natural disasters. Therefore, this study also provided a unique opportunity to examine the use of Twitter to detect dietary patterns.

The present study examined mentions of various foods commonly high in energy, protein, fat, and/or carbohydrate on Twitter that were posted in conjunction with posts related to hurricanes. Results of this study indicated that foods mentioned on Twitter during four hurricanes were commonly energy and protein dense. While the present study did not find any differences in food group mentions overall or post-hurricane, there were significant differences in food groups mentioned pre- and during the hurricanes. In particular, foods categorized as grains were commonly mentioned before the hurricanes and foods categorized as dairy (mainly milk) were most commonly mentioned during the hurricanes. It is not surprising the milk and grain-based foods (particularly breads and rolls) were commonly tweeted foods before and during the hurricanes. Stockpiling milk, along with bread and eggs, is commonly reported in the media in advance of storms (CoxMedia 2018). This may be one of the reasons why milk and bread were so frequently mentioned. Additionally, pizza, which is essentially a combination of grains and dairy, was frequently mentioned over most of the time points for the hurricanes. The previous research has shown that pre-hurricane is a time when individuals may be attempting to use up perishable foods they

have in their household (Pyles et al. 2008). In addition, individuals may be purchasing non-perishable forms of foods that would usually require refrigeration (“canned milk”) (Pyles et al. 2008) or purchasing cooking supplies that do not rely on electricity (charcoal grills) (Pyles et al. 2008).

While mentions of milk and bread seem to align with previous reports of food purchases prior to natural disasters, mentions of other foods were surprising. For example, apples and oranges were frequently mentioned. It is possible that people were tweeting about the color orange or Orange County, Florida, rather than the actual fruit. However, the results clearly mirrored the findings that oranges and apples are the two most commonly consumed fruits in the USA, although mostly consumed in the form of juice (USDA 2019). In addition, peanuts were frequently mentioned, particularly after the hurricanes. It is unclear whether peanuts are a common, non-perishable food people may have on hand to consume or the mentions referred to something else, such as categorizing the size of hail during a storm (Weather.gov 2019). For meat, white meat and poultry were most frequently cited as compared to red meat, such as hamburgers, sausage, or pork. While seafood consumption has decreased in the USA (Terry et al. 2018), poultry intake has been on the rise, surpassing beef intake (USDA 2019). Lastly, of the four common beverages mentioned, three were alcoholic beverages (beer, alcohol, and wine). While disaster preparation guidelines recommend avoiding alcohol and focusing on healthy, non-perishable foods (CDC 2014), the findings of this analysis show that many people indicated that they were not following this guideline when posting to Twitter. While alcohol was frequently mentioned, this may be more of a reflection of the common use of social media to discuss drinking, especially among young adults (Hebden et al. 2015; Lyons et al. 2015), than a reflection of a disproportionate intake of alcohol over other F&B groups. For example, users of social media may be more likely to post about alcohol use versus consumption of legumes.

Of all the food groups assessed in this study, vegetables had the fewest mentions. This is possibly because these items are typically perishable. However, meat and dairy foods were frequently mentioned, and they can also be perishable. It is likely that vegetables were not frequently mentioned because intakes of vegetables in the USA are very low and usually only in the form of French fries (potatoes) or pizza sauce (tomatoes) (USDA 2019). Vegetable intake is particularly low in the southern US, as compared to other regions (PBH 2015). Foods categorized as desserts and sweets were also not frequently mentioned. This is surprising given the fact that refined sugar intake has been increasing (USDA 2019) and that snack foods, such as cookies and cakes, make up to 25% of US adults’ energy intake (Dunford and Popkin 2017). It is possible that these foods are viewed as not necessary for stockpiling in preparation of a storm. Despite the possibility that individuals susceptible to emotional eating would be at risk for increasing intake of sweets after a natural disaster, the previous research found that there was not an increase in junk foods (both savory and sweet snacks) after an earthquake among women with emotional eating (Kuijer and Boyce 2012). Findings of that study did indicate that stress in response to natural disasters can lead to a decrease in intake of healthy foods, such as vegetables (Kuijer and Boyce 2012). This may explain the low reported mentions of vegetables in the present study.

Overall, foods mentioned in the present study were high in energy and protein. Findings indicated that efforts should be made to encourage individuals preparing for a storm to purchase healthy foods to have on hand. Considering that foods high in protein, such as poultry, dairy, eggs, and seafood, are perishable and are also some of the top contributors to food-borne illness (CDC 2018), efforts to educate consumers on shelf-stable sources of protein, including peanut butter and canned legumes, may be warranted. The CDC, as well as state and local agencies, may wish to increase education efforts around having healthy,

non-perishable food on hand before storms arrive. The CDC has used viral social media campaigns to increase preparedness for natural disasters, but there has been no previous concentrated focus on healthy foods to have on hand for disasters (Fraustino and Ma 2015).

The present findings should be considered in the context of some potential limitations. It is possible that some of the F&B mentions contained names of foods, but the user could have been tweeting about something not food related (e.g., “This weather is nuts!”). In addition, only a sampling of approximately 500,000 tweets containing the hurricanes hashtag names or full names were collected. Some foods were mixed dishes, such as waffles and biscuits (which may contain grains, dairy, and eggs). In addition, while pizza was categorized as a grain, it is one of the largest contributors to dairy consumption in the USA, with Italian cheeses being consumed more frequently than any other cheese (USDA 2014). A broad range of foods and beverages were selected based on known energy and macronutrient contributors in the USA (O’Neil et al. 2012); however, it could also be that, for example, people tweeted about Oreos™ instead of “cookies.” While a quarter of US adults use Twitter, users of the platform may not be generalizable to the US population, with the majority of Twitter users being under the age of 50 (67%) (Pew 2018) and tending to be more highly educated and wealthier than non-Twitter users (Wojcik and Hughes 2019). However, there is equal distribution among users by sex, race, education, and urban/suburban locations (Pew 2018). While the previous research has shown that the most commonly tweeted foods by US region align with traditional regional cuisine (e.g., grits are the most commonly tweeted food word in the southeastern US) (Fried et al. 2014), it is possible that people may be more likely to tweet about unusual foods versus foods they consume every day. Lastly, actual consumption was not measured; only mentions of F&B. Future studies may wish to employ a mixed methods approach to examining the content of messages. For example, a more thematic or content analysis approach could be conducted, including an examination of images that are posted to Twitter. An analysis of images could help provide further context of the messages posted to Twitter.

The study also has several strengths. Using a framework of known major contributors to energy and macronutrients allowed for the collection of individual foods and beverages, as well as an examination of dietary patterns by energy and macronutrients. The methods used in this study could also be applied to other major events as a way to study dietary patterns. This includes during other natural disasters (e.g., winter storms), sporting events (e.g., the Super Bowl), and variations in seasons (e.g., eating patterns in summer vs. winter). The resulting list of F&B and broader food group categories could be used for other studies and with other social media platforms. The methods could also be applied to other “big data” platforms, such as other social media sites, electronic health records, and diet tracking mobile apps.

5 Conclusion

Social media may be a unique way to detect dietary patterns and help inform recommendations for what F&B are appropriate during natural disasters. The present study found frequent mentions of grains and dairy, particularly pizza and milk, as well as foods that are energy dense and high in protein. Since healthy food items, such as fruits and vegetables, were infrequently mentioned, officials in charge of making disaster preparedness recommendations related to food may want to encourage the purchase of fruits (such as bananas or dried fruits) and vegetables (such as canned peas or carrots) that can be stored

and consumed without the need for electricity. Disaster preparedness officials may also want to reinforce the message about limiting or avoiding alcohol before, during, and after a natural disaster.

Funding This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Bennett DM (2014) How do emergency managers use social media platforms? *J Emerg Manag* 12(3):251–256. <https://doi.org/10.5055/jem.2014.0178>
- Brandt HM, Turner-McGrievy G, Friedman DB, Gentile D, Schrock C, Thomas T, West D (2019) Examining the role of twitter in response and recovery during and after historic flooding in South Carolina. *J Public Health Manag Pract* 25(5):E6–E12. <https://doi.org/10.1097/phh.0000000000000841>
- CDC (2014) Centers for disease control and prevention. Crisis and Emergency Risk Communication 2014 edition. https://emergency.cdc.gov/cerc/resources/pdf/cerc_2014edition.pdf. Accessed 20 Mar 2019
- CDC (2017) Center for disease control and prevention. Before a hurricane. <https://www.cdc.gov/disasters/hurricanes/before.html>. Accessed 20 Mar 2019
- CDC (2018) Foods linked to food poisoning. Centers for Disease Control and Prevention, National Center for Emerging and Zoonotic Infectious Diseases (NCEZID), Division of Foodborne, April 19, 2018
- CDC (2019) Centers for disease control and prevention. Food and Water Needs: Preparing for a Disaster or Emergency. <https://www.cdc.gov/disasters/foodwater/prepare.html>. Accessed 20 Mar 2019
- CoxMedia(2018) Cox media group national content desk. Bread and milk: why do we panic-buy those foods before a storm? The Atlanta Journal-Constitution. May 16, 2018. <https://www.ajc.com/weather/bread-and-milk-why-panic-buy-those-foods-before-storm/QJoUPzPhRMpn4Gx5naSuYP/>. Accessed 1 Apr 2019
- Dedaj P (2017) Drinking at ‘hurricane parties’ a bad idea, Florida officials warn. Fox News. <https://www.foxnews.com/us/drinking-at-hurricane-parties-a-bad-idea-florida-officials-warn>. Accessed 20 Mar 2019
- Der-Martirosian C, Strine T, Atia M, Chu K, Mitchell MN, Dobalian A (2014) General household emergency preparedness: a comparison between veterans and nonveterans. *Prehosp Disaster Med* 29(2):134–140. <https://doi.org/10.1017/s1049023x1400020x>
- Duncan C (2019) News and observer. Hurricane Florence was among the costliest disasters on record. Here’s NOAA’s tally. March 4, 2019. <https://www.newsobserver.com/news/state/north-carolina/article225974185.html>. Accessed 21 Mar 2019
- Dunford EK, Popkin BM (2017) Disparities in snacking trends in US Adults over a 35 year period from 1977 to 2012. *Nutrients* 9(8):809
- Finch KC, Snook KR, Duke CH, Fu K-W, Tse ZTH, Adhikari A, Fung IC-H (2016) Public health implications of social media use during natural disasters, environmental disasters, and other environmental concerns. *Nat Hazards* 83(1):729–760. <https://doi.org/10.1007/s11069-016-2327-8>
- Flory K, Hankin BL, Kloos B, Cheely C, Turecki G (2009) Alcohol and cigarette use and misuse among Hurricane Katrina survivors: psychosocial risk and protective factors. *Subst Use Misuse* 44(12):1711–1724. <https://doi.org/10.3109/10826080902962128>
- Fraustino JD, Ma L (2015) CDC’s use of social media and humor in a risk campaign—“preparedness 101: Zombie Apocalypse”. *J Appl Commun Res* 43(2):222–241. <https://doi.org/10.1080/00909882.2015.1019544>
- Fried D, Surdeanu M, Kobourov S, Hingle M, Bell D (2014) Analyzing the language of food on social media. Paper presented at the 2014 IEEE international conference on big data (big data)
- Gore RJ, Diallo S, Padilla J (2015) You are what you Tweet: connecting the geographic variation in America’s obesity rate to Twitter content. *PLoS ONE* 10(9):e0133505. <https://doi.org/10.1371/journal.pone.0133505>
- Guan X, Chen C (2014) Using social media data to understand and assess disasters. *Nat Hazards* 74(2):837–850. <https://doi.org/10.1007/s11069-014-1217-1>
- Guskin E, Hirtline P (2012) Hurricane Sandy and Twitter. <http://www.journalism.org/2012/11/06/hurricane-sandy-and-twitter/>. Accessed 10/12/17

- Harris JK, Hawkins JB, Nguyen L, Nsoesie EO, Tuli G, Mansour R, Brownstein JS (2017) Using Twitter to identify and respond to food poisoning: the food safety STL project. *J Public Health Manag Pract* 23(6):577–580. <https://doi.org/10.1097/phh.0000000000000516>
- Hebden R, Lyons AC, Goodwin I, McCreanor T (2015) “When you add alcohol, it gets that much better”: university students, alcohol consumption, and online drinking cultures. *J Drug Issues* 45(2):214–226. <https://doi.org/10.1177/0022042615575375>
- Heimerl F, Lohmann S, Lange S, Ertl T (2014) Word cloud explorer: text analytics based on word clouds. Paper presented at the 2014 47th Hawaii international conference on system sciences
- Heimlich R (2012) Pew research center. Twitter a Key Source of news for many during Hurricane Sandy. November 14, 2012. <http://www.pewresearch.org/fact-tank/2012/11/14/twitter-a-key-source-of-news-for-many-during-hurricane-sandy/>. Accessed 20 Mar 2019
- Hingle M, Yoon D, Fowler J, Kobourov S, Schneider ML, Falk D, Burd R (2013) Collection and visualization of dietary behavior and reasons for eating using Twitter. *J Med Internet Res* 15(6):e125. <https://doi.org/10.2196/jmir.2613>
- Karami A, Dahl AA, Turner-McGrievy G, Kharrazi H, Shaw G Jr (2018) Characterizing diabetes, diet, exercise, and obesity comments on Twitter. *Int J Inf Manag* 38(1):1–6
- Karami A, Shah V, Vaezi R, Bansal A (2019) Twitter speaks: a case of national disaster situational awareness. *J Inf Sci*. <https://doi.org/10.1177/0165551519828620>
- Karami A, Lundy M, Webb F, Dwivedi YK (2020) Twitter and research: a systematic literature review through text mining. *IEEE Access* 8:67698–67717
- Kuijter RG, Boyce JA (2012) Emotional eating and its effect on eating behaviour after a natural disaster. *Appetite* 58(3):936–939. <https://doi.org/10.1016/j.appet.2012.02.046>
- Kuttschreuter M, Hilverda F (2019) “Listen, did you hear...?” A structural equation model explaining online information sharing on the risks of nanotechnology in food. *Food Qual Prefer*. <https://doi.org/10.1016/j.foodqual.2019.03.011>
- Laseter E (2018) These are the best healthy foods to stockpile for a Hurricane. *Cooking Light*. September 10, 2018. <https://www.cookinglight.com/cooking-101/healthy-hurricane-food-tips>. Accessed 20 Mar 2019
- Lyons AC, Goodwin I, McCreanor T, Griffin C (2015) Social networking and young adults’ drinking practices: innovative qualitative methods for health behavior research. *Health Psychol* 34(4):293–302. <https://doi.org/10.1037/hea0000168>
- Nguyen QC, Meng H, Li D, Kath S, McCullough M, Paul D, Li F (2017) Social media indicators of the food environment and state health outcomes. *Public Health* 148:120–128. <https://doi.org/10.1016/j.puhe.2017.03.013>
- O’Neil CE, Keast DR, Fulgoni VL, Nicklas TA (2012) Food sources of energy and nutrients among adults in the US: NHANES 2003–2006. *Nutrients* 4(12):2097–2120. <https://doi.org/10.3390/nu4122097>
- PBH, Produce for Better Health Foundation (2015) State of the plate: 2015 study on America’s consumption of fruit & vegetables
- Pew (2018) Pew Research Center: Internet & Technology. Social Media Fact Sheet. February 5, 2018. <https://www.pewinternet.org/fact-sheet/social-media/>. Accessed 1 Apr 2019
- Pyles L, Kulkarni S, Lein L (2008) Economic survival strategies and food insecurity. *J Soc Serv Res* 34(3):43–53. <https://doi.org/10.1080/01488370802086047>
- RCoreTeam (2014) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. <http://www.R-project.org/>
- Scott KK, Errett NA (2018) Content, accessibility, and dissemination of disaster information via social media during the 2016 Louisiana floods. *J Public Health Manag Pract* 24(4):370–379. <https://doi.org/10.1097/phh.0000000000000708>
- Sinnenberg L, Bittenheim AM, Padrez K, Mancheno C, Ungar L, Merchant RM (2017) Twitter as a tool for health research: a systematic review. *Am J Public Health* 107(1):e1–e8. <https://doi.org/10.2105/ajph.2016.303512>
- Smith K (2019) 122 amazing social media statistics and facts. <https://www.brandwatch.com/blog/amazing-social-media-statistics-and-facts/>. Accessed 2/21/19
- Smith A, Anderson M (2018) Social media use in 2018. Pew Research Center. <http://www.pewinternet.org/2018/03/01/social-media-use-in-2018/>
- Sutton J, League C, Sellnow TL, Sellnow DD (2015) Terse messaging and public health in the midst of natural disasters: the case of the Boulder floods. *Health Commun* 30(2):135–143. <https://doi.org/10.1080/10410236.2014.974124>
- Terry AL, Herrick KA, Afful J, Ahluwalia N (2018) Seafood consumption in the United States, 2013–2016. NCHS data brief no. 321, pp 1–8

- USDA, United States Department of Agriculture (2014) Statistic: food consumption & demand. Trends in U.S. per capita consumption of dairy products, 1970–2012. <https://www.ers.usda.gov/amber-waves/2014/june/trends-in-us-per-capita-consumption-of-dairy-products-1970-2012/>
- USDA, United States Department of Agriculture (2019) Food availability and consumption. <https://www.ers.usda.gov/data-products/ag-and-food-statistics-charting-the-essentials/food-availability-and-consumption/>
- Vidal L, Ares G, Machín L, Jaeger SR (2015) Using Twitter data for food-related consumer research: a case study on “what people say when tweeting about different eating situations”. *Food Qual Prefer* 45:58–69. <https://doi.org/10.1016/j.foodqual.2015.05.006>
- Wang C (2017) Bracing for Hurricanes: a qualitative analysis of the extent and level of preparedness among older adults. *Gerontologist* 58(1):57–67. <https://doi.org/10.1093/geront/gnx187>
- Wang B, Zhuang J (2017) Crisis information distribution on Twitter: a content analysis of tweets during Hurricane Sandy. *Nat Hazards* 89(1):161–181. <https://doi.org/10.1007/s11069-017-2960-x>
- Wang B, Zhuang J (2018) Rumor response, debunking response, and decision makings of misinformed Twitter users during disasters. *Nat Hazards* 93(3):1145–1162. <https://doi.org/10.1007/s11069-018-3344-6>
- Wang Z, Ye X, Tsou M-H (2016) Spatial, temporal, and content analysis of Twitter for wildfire hazards. *Nat Hazards* 83(1):523–540. <https://doi.org/10.1007/s11069-016-2329-6>
- Weather.gov (2019) National weather service: estimating hail size. <https://www.weather.gov/boi/hailsize>
- Wojcik S, Hughes A (2019) Sizing up Twitter users. April 24, 2019. <https://www.pewinternet.org/2019/04/24/sizing-up-twitter-users/>. Accessed 10-4-19
- Xiao Y, Huang Q, Wu K (2015) Understanding social media data for disaster management. *Nat Hazards* 79(3):1663–1679. <https://doi.org/10.1007/s11069-015-1918-0>
- Zhou J, Liu F, Zhou H (2018) Understanding health food messages on Twitter for health literacy promotion. *Perspect Public Health* 138(3):173–179. <https://doi.org/10.1177/1757913918760359>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.